

Supplementary Table 1. STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Section & paragraph No.
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Abstract
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Abstract
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Introduction 1, 2, 3
Objectives	3	State specific objectives, including any prespecified hypotheses	Introduction 4
Methods			
Study design	4	Present key elements of study design early in the paper	Methods, Study population 1
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Method, Study population 1, 2
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	Method, Study population 1, 2
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	NA
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Methods, Variables and Outcomes 1
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Methods, Variables and Outcomes 1
Bias	9	Describe any efforts to address potential sources of bias	NA
Study size	10	Explain how the study size was arrived at	Method, Patients and Samples 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Method, Statistical Methods 1
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Method, Statistical Methods 1
		(b) Describe any methods used to examine subgroups and interactions	Method, Statistical Methods 1
		(c) Explain how missing data were addressed	NA
		(d) Cohort study—If applicable, explain how loss to follow-up was addressed Case-control study—If applicable, explain how matching of cases and controls was addressed Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy	Method, Statistical Methods 1
		(e) Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—e.g., numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Results, Baseline characteristics of study population 1, 2
		(b) Give reasons for non-participation at each stage	NA
		(c) Consider use of a flow diagram	NA

Supplementary Table 1. Continued

	Item No	Recommendation	Section & paragraph No.
Descriptive data	14*	(a) Give characteristics of study participants (e.g., demographic, clinical, social) and information on exposures and potential confounders	Table 1 Results, Baseline characteristics of study population 1, 2
		(b) Indicate number of participants with missing data for each variable of interest	NA
		(c) Cohort study—Summarise follow-up time (e.g., average and total amount)	NA
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time	Figure 1A-1D Results, Overall survival according to etiologies 1
		Case-control study—Report numbers in each exposure category, or summary measures of exposure	NA
		Cross-sectional study—Report numbers of outcome events or summary measures	NA
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Results, Predictive factors for survival in patients with HCC 1
		(b) Report category boundaries when continuous variables were categorized	NA
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	Results, The effects of diabetes mellitus on the risk of death 1
Discussion			
Key results	18	Summarise key results with reference to study objectives	Discussion 1
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Discussion 3
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Discussion 2,3,4
Generalisability	21	Discuss the generalisability (external validity) of the study results	Discussion 4
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Funding statement

An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.